


arkray

Urinalysis reagent for visual or machine reading

# AUTION Sticks

## AUTION Sticks 10PA



Creatinine correction is vital for protein measurement with single voided urine.


Conventional urinary tests using single voided urine sometimes undervalue or overvalue urine protein due to increased or reduced urine concentration.

# High-precision screening of kidney disease is now possible even with single-voided urine.

## Interpretation of Test Results

Glu					
Semiquantitative symbol	NORMAL	±	+1	+2	+3
Conc. (mg/dL)		50	100	200	500
Pro					
Semiquantitative symbol	NEG.	±	+1	+2	+3
Conc. (mg/dL)		15	30	100	300
Bil					
Semiquantitative symbol	NEG.	+1	+2	+3	+4
Conc. (mg/dL)		0.5	2	6	OVER
Uro					
Semiquantitative symbol	NORMAL	+1	+2	+3	+4
Conc. (mg/dL)		2	4	8	OVER
pH		5.0	6.0	7.0	8.0
Bld					
Semiquantitative symbol	NEG.	+1	+2	+3	
Conc. (mg/dL)		0.06	0.2	1.0	
Ket					
Semiquantitative symbol	NEG.	±	+1	+2	+3
Conc. (mg/dL)		NORMAL	15	40	80
Nit					
Semiquantitative symbol	NEG.	+1	+2		
Leu					
Conc. (Leu/μL)	NEG.	25	75	250	500
Cre					
Conc. (mg/dL)		10	50	100	200
					300

## Correlation between P/C ratio and A/C ratio in concentration



## Protein / Creatinine ratio

Amount required for judgement	Creatinine (mg/dL)				
	10	50	100	200	300
NEG.	Re-analysis	normal	normal	normal	normal
15	+2	+1	+1	normal	normal
30	+2	+2	+1	+1	+1
100	+2	+2	+2	+2	+1
300	+2	+2	+2	+2	+2
1000	+2	+2	+2	+2	+2

## Specification

Measurement sample	Urine (fresh urine, voided urine)
Measurement item	Glu, Pro, Bil, pH, Bld, Uro, Ket, Nit, Leu, Cre, (Calculated item/ Pro/Cre comparison)
Required sample volume	Approx. 5mL (Dipping method)
Reaction time	60 seconds (Leu:90 sec.)
Reaction temperature	Room temperature
Preservation conditions	At room temperature Avoid direct sunlight
Expiry	24 months


## Reaction principle

Test item	Reaction principle
Glucose	Glucose oxidase method
Protein	Protein-error method
Bilirubin	Azo-coupling method
Urobilinogen	Azo-coupling method
pH	pH indicator method
Blood	Activity measurement of pseudoperoxidase in haemoglobin
Ketones	Sodium nitroprusside method
Nitrite	Greiss method
Leucocyte	Leucocyte esterase activity method
Creatinine	Chelate competition method

## Creatinine

When Creatinine pigment in the urine reacts with palladium compound, blue-green color on test paper changes to yellow. The decreased ratio of blue-green color level is proportional to creatinine concentration.

### Competition reaction



ARKRAY GLOBAL BUSINESS, INC.

KYOTO MIYUKI Bldg. 10F, 689 Takanna-cho,  
Nakagyo-ku, Kyoto 604-8153, JAPAN

TEL +81-75-662-8967 FAX +81-75-662-8973

